



DO NOT OPEN

UNTIL INSTRUCTED TO DO SO

CHEM 110 – Dr. McCorkle – Exam #3

While you wait, please complete the following information:

Name: _____

Student ID: _____

Turn off cellphones and stow them away. No headphones, mp3 players, hats, sunglasses, food, drinks, restroom breaks, graphing calculators, programmable calculators, or sharing calculators. Grade corrections for incorrectly marked or incompletely erased answers will not be made.

Periodic Table of the Elements

PERIOD	GROUP 1 IA	2 IIA	3 IIIB	4 IVB	5 VB	6 VIB	7 VIIB	8 VIII	9 VIII	10 VIII	11 IB	12 IB	13 IIIA	14 IVA	15 VA	16 VIA	17 VIIA	18 VIIIA
1	1 H 1.01	2 He 4.00																
2	3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
3	11 Na 22.99	12 Mg 24.31											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95
4	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.88	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.61	33 As 74.92	34 Se 78.97	35 Br 79.90	36 Kr 83.80
5	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.95	43 Tc (98)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.75	52 Te 127.60	53 I 126.90	54 Xe 131.29
6	55 Cs 132.91	56 Ba 137.33	57 La* 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.85	75 Re 186.21	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.2	83 Bi 208.98	84 Po (209)	85 At (210)	86 Rn (222)
7	87 Fr (223)	88 Ra (226)	89 Ac** (227)	104 Rf (267)	105 Db (268)	106 Sg (271)	107 Bh (270)	108 Hs (277)	109 Mt (276)	110 Ds (281)	111 Rg (280)	112 Cn (285)	113 Uut (284)	114 Fl (289)	115 Uup (288)	116 Lv (293)	117 Uus (294)	118 Uuo (294)

Lanthanide Series *

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.12	140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.05	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(262)

Actinide Series **

Multiple Choice – Choose the answer that best completes the question. Use an 815-E Scantron to record your response. [2 points each]

- In the kinetic molecular theory of gas behavior, the assumption is made that gas molecules
 - occasionally come to rest.
 - are attracted to each other by strong forces.
 - are close together in their container.
 - move with a kinetic energy equal to their centigrade temperature.
 - move rapidly in random directions.
- The atmospheric pressure in Denver, CO is 633 mmHg. What is this pressure in atm?
 - 1.20 atm
 - 633 atm
 - 0.833 atm
 - 1.00 atm
 - 127 atm
- The volume of a gas with a pressure of 1.2 atm increases from 1.0 L to 4.0 L. What is the final pressure of the gas, assuming no change in moles or temperature?
 - 1.2 atm
 - 0.30 atm
 - 3.3 atm
 - 4.8 atm
 - 1.0 atm
- Complete the following statement: In Charles's Law, the volume of a gas _____ when the _____ decreases.
 - increases; temperature
 - increases; quantity of gas
 - increases; pressure
 - decreases; temperature
 - decreases; pressure
- A gas at 5.00 atm pressure was stored in a tank during the winter at 5.0 °C. During the summer, the temperature in the storage area reached 40.0 °C. What was the pressure in the gas tank then?
 - 0.625 atm
 - 4.44 atm
 - 5.63 atm
 - 40.0 atm
 - 69.5 atm
- A mixture of 10.0 g of Ne and 10.0 g of Ar have a total pressure of 1.6 atm. What is the partial pressure of Ne?
 - 0.40 atm
 - 0.54 atm
 - 0.80 atm
 - 1.1 atm
 - 1.3 atm
- How many moles of neon occupy a volume of 14.3 L at STP?
 - 36.7 moles
 - 32.0 moles
 - 6.45 moles
 - 0.638 moles
 - 1.57 moles
- Calculate the root mean square velocity of nitrogen molecules at 25 °C.
 - 149 m/s
 - 297 m/s
 - 515 m/s
 - 729 m/s
 - 1090 m/s


9. Define *energy*.
- A) the flow of energy caused by a chemical reaction
 - B) the flow of energy caused by a temperature difference
 - C) the result of a force acting through a distance
 - D) the capacity to do work
 - E) a chemical reaction
10. The gas in a piston (defined as the system) is warmed and absorbs 640. J of heat. The expansion performs 344 J of work on the surroundings. What is the change in internal energy for the system in joules?
- A) 296 J B) 984 J C) -296 J D) -984 J E) 1.86 J
11. Which is the correct equation for the formation of $\text{Na}_2\text{CO}_3(s)$ from its elements in their standard states?
- A) $2 \text{Na}(s) + \text{C}(s, \textit{graphite}) + 3 \text{O}(g) \rightarrow \text{Na}_2\text{CO}_3(s)$
 - B) $2 \text{Na}(s) + \text{C}(s, \textit{graphite}) + 3/2 \text{O}_2(g) \rightarrow \text{Na}_2\text{CO}_3(s)$
 - C) $2 \text{Na}^+(aq) + \text{CO}_3^{2-}(aq) \rightarrow \text{Na}_2\text{CO}_3(s)$
 - D) $4 \text{Na}(s) + 2 \text{C}(s, \textit{graphite}) + 6 \text{O}(g) \rightarrow 2 \text{Na}_2\text{CO}_3(s)$
 - E) $4 \text{Na}(s) + 2 \text{C}(s, \textit{graphite}) + 3 \text{O}_2(g) \rightarrow 2 \text{Na}_2\text{CO}_3(s)$

Calculations – Write your initials in the upper-right corner of every page that contains work. For full credit show all work and write neatly; give answers with correct significant figures and units. Place a box around your final answer.

12. A sample of nitrogen gas had a volume of 500. mL, a pressure in its closed container of 640. torr, and a temperature of 25 °C. What was the new volume of the gas in liters when the temperature was changed to 50 °C and the new pressure was 1.00 atm? [6 points]

13. A 0.334 g sample of an unknown halogen occupies 109 mL at 125 °C and 1.41 atm. What is the identity of the halogen? [5 points]


14. Consider the following reaction:



When 35.0 g of $\text{Al}_2(\text{CO}_3)_3$ reacts, how many L of CO_2 gas are formed at 55 °C and a pressure of 975 mmHg? [5 points]


15. A 4.98 g sample of aniline ($\text{C}_6\text{H}_5\text{NH}_2$) was combusted in a bomb calorimeter with a heat capacity of 4.25 kJ/°C. If the temperature rose from 29.5 °C to 69.8 °C, determine the value of ΔE_{rxn} for aniline. [4 points]

16. What volume in mL of benzene (C_6H_6 , $d = 0.88 \text{ g/mL}$) is required to produce $1.5 \times 10^3 \text{ kJ}$ of heat according to the following reaction: [4 points]





17. A piece of iron (mass = 25.0 g) at 125 °C is placed in a Styrofoam coffee cup containing 25.0 mL of water at 20. °C. Assuming that no heat is lost to the cup or the surroundings, what will the final temperature of the water be? ($d_{\text{water}} = 0.997 \text{ g/mL}$, $s_{\text{Fe}} = 0.449 \text{ J/g}\cdot^\circ\text{C}$, $s_{\text{water}} = 4.184 \text{ J/g}\cdot^\circ\text{C}$) [5 points]


18. Calculate ΔH_{rxn} for the following reaction: [4 points]



Use the following reactions and given ΔH 's.



19. Calculate $\Delta H_{\text{f}}^{\circ}$ of $\text{SO}_2(g)$ using the following reaction and standard enthalpies of formation:
[5 points]



20. Acetylene gas (C_2H_2) is commonly used in welding torches. When burned with pure oxygen, instead of air, it burns at about $3,500\text{ }^\circ\text{C}$!

a. Write a balanced equation for the combustion of acetylene (C_2H_2) in pure oxygen. (Products are gaseous due to the high temperature.) [2 points]

b. Calculate the $\Delta H^\circ_{\text{rxn}}$ using the standard enthalpies of formation. [4 points]

c. Two pieces of iron at $23\text{ }^\circ\text{C}$ have a total mass of 1250 g. What is the final temperature of the iron in Celsius if 25.5 g of acetylene is used to weld them together? Assume all of the heat from the combustion of acetylene is transferred to the iron. ($s_{\text{Fe}} = 0.449\text{ J/g}\cdot^\circ\text{C}$) [6 points]

21. The tiles on the space shuttle have an incredible specific heat capacity of $62.8 \text{ J/g}\cdot^\circ\text{C}$ and were designed to withstand incredible temperatures. Let's imagine that after a successful mission, the space shuttle reenters Earth's atmosphere where its external temperature rises from -157°C to 1650°C generating $5.38 \times 10^{11} \text{ J}$ of heat. If the average mass of each tile is 195 g , how many tiles would you need to absorb this heat? (Assume the heat is distributed evenly among the tiles.) [5 points]

Extra Credit: What Italian scientist invented the barometer? [2 points]

**Formulas & Constants
(you may or may not need)**

1 inch = 2.54 cm (exact)

1 mile = 5280 ft \approx 1.609 km

1 kg \approx 2.205 lb

1 lb = 16 oz \approx 453.6 g

1 gal = 4 qt = 8 pt \approx 3.785 L

1 L = 1000 cm³

K = °C + 273.15

°F = 1.8 x °C + 32

°C = (°F – 32)/1.8

1 cal = 4.184 J

1 Cal = 1000 cal

q = m x C x Δ T

Avogadro's # = 6.022x10²³

Molar volume = 22.4 L

R = 0.08206 $\frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}}$

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

$$u_{\text{rms}} = \sqrt{\frac{3RT}{M}}$$

$$KE = \frac{1}{2}mv^2 = \frac{3}{2}RT$$

1 atm = 760 mmHg

1 mmHg = 1 torr

P_{Total} = P₁ + P₂ + ...

P_A = X_A·P_{Total}

PV = nRT

$\Delta E = q + w$

w = -P Δ V

q = C x Δ T

q = m x s x Δ T

$$\Delta H^\circ_{\text{rxn}} = \Sigma[n \Delta H^\circ(\text{products})] - \Sigma[n \Delta H^\circ(\text{reactants})]$$

R = 8.314 J/mol·K

Standard Enthalpies of Formation at 25 °C

Substance	ΔH°_f (kJ/mol)
CH ₄ (g)	-74.9
C ₂ H ₂ (g)	+227.4
C ₆ H ₆ (l)	+49.1
CO ₂ (aq)	-413.8
CO ₂ (g)	-393.5
H ₂ O(g)	-241.8
H ₂ O(l)	-285.8
H ₂ S(aq)	-39.4
H ₂ S(g)	-20.6

Scratch Page
(to be handed in)