



DO NOT OPEN
UNTIL INSTRUCTED TO DO SO

CHEM 110 – Dr. McCorkle – Exam #4

While you wait, please complete the following information:

Name: _____

Student ID: _____

Turn off cellphones and stow them away. No headphones, mp3 players, hats, sunglasses, food, drinks, restroom breaks, graphing calculators, programmable calculators, or sharing calculators. Grade corrections for incorrectly marked or incompletely erased answers will not be made.

GROUP 1 | A

PERIOD

Lanthanide Series*

Actinide Series**

Multiple Choice – Choose the answer that best completes the question. Use an 815-E Scantron to record your response. [2 points each]

1. The vertical height of a wave is called
 - A) wavelength
 - B) amplitude
 - C) frequency
 - D) energy
 - E) de Broglie wavelength
2. When waves of equal amplitude from two sources are out of phase when they interact, it is called
 - A) destructive interference
 - B) diffraction
 - C) amplitude
 - D) constructive interference
 - E) uncertainty
3. Which of the following quantum numbers describes the shape of an orbital?
 - A) principal quantum number
 - B) magnetic quantum number
 - C) spin quantum number
 - D) Schrödinger quantum number
 - E) angular momentum quantum number
4. Which of the following quantum numbers describes the orientation of an orbital?
 - A) principal quantum number
 - B) magnetic quantum number
 - C) spin quantum number
 - D) Schrödinger quantum number
 - E) angular momentum quantum number
5. Electrons filling up the orbitals from low energy to high energy is known as (the)
 - A) Pauli exclusion principle
 - B) Hund's rule
 - C) Aufbau principle
 - D) Heisenberg uncertainty principle
 - E) Coulomb's law

6. How many valence electrons do the halogens possess?
A) 1 B) 2 C) 5 D) 6 E) 7
7. Suppose a metal ejects electrons from its surface when struck by red light. What will happen if the surface is struck by blue light?
A) No electrons would be ejected.
B) Electrons would be ejected, and they would have the same kinetic energy as those ejected by red light.
C) Electrons would be ejected, and they would have greater kinetic energy than those ejected by red light.
D) Electrons would be ejected, and they would have lower kinetic energy than those ejected by red light.
8. Identify the isoelectronic elements
A) Cl^- , F^- , Br^- , I^- , At^-
B) N^{3-} , S^{2-} , Br^- , Cs^+ , Sr^{2+}
C) P^{3-} , S^{2-} , Cl^- , K^+ , Ca^{2+}
D) Zn^{2+} , Co^{2+} , Cu^{2+} , Cr^{2+} , Cd^{2+}
E) Ne, Ar, Kr, Xe, He
9. What period 3 element has the following ionization energies (all in kJ/mol)?
 $\text{IE}_1 = 1012$ $\text{IE}_2 = 1900$ $\text{IE}_3 = 2910$ $\text{IE}_4 = 4960$ $\text{IE}_5 = 6270$ $\text{IE}_6 = 22,200$
A) Mg B) Si C) P D) S E) Cl
10. Arrange the elements in order of increasing IE_1 : N, F, As
A) $\text{N} < \text{As} < \text{F}$
B) $\text{As} < \text{N} < \text{F}$
C) $\text{F} < \text{N} < \text{As}$
D) $\text{As} < \text{F} < \text{N}$
E) $\text{F} < \text{As} < \text{N}$
11. Arrange the elements in order of increasing metallic character: P, As, K
A) $\text{P} < \text{As} < \text{K}$
B) $\text{As} < \text{P} < \text{K}$
C) $\text{K} < \text{P} < \text{As}$
D) $\text{As} < \text{K} < \text{P}$
E) $\text{K} < \text{As} < \text{P}$

Calculations – Write your initials in the upper-right corner of every page that contains work. For full credit show all work and write neatly; give answers with correct significant figures and units. For calculations, place a box around your final answer.

12. For the following short answer questions, please circle, fill-in, or provide the correct answer.

[2 points each]

- a. Which transition has a longer wavelength?

$$n = 7 \rightarrow n = 5$$

$$n = 3 \rightarrow n = 1$$

- b. Which has the lowest energy?

6s

5d

4f

- c. A gas phase element gaining an electron is a typically endothermic process.

True

False

- d. How many electrons in an atom could have $n = 4$, $l = 2$, $m_l = +1$? _____

- e. The two most probable ions of titanium are _____ and _____.

13. How many photons are contained in a flash of green light (525 nm) that contains 189 kJ of energy? [4 points]

14. Determine which of the following represents an acceptable set of quantum numbers for an electron in an atom. If the set is acceptable then circle the entire row. If it is not then draw a line through the entire row. [2 points each]

a. $n = 4$ $l = 2$ $m_l = -1$

b. $n = 3$ $l = 1$ $m_l = +2$

c. $n = 5$ $l = 0$ $m_l = 0$

15. Complete the following table: [9 points]

Symbol	Condensed Electron Configuration	# of Valence e ⁻	# of Unpaired e ⁻
Bi			
Co ⁴⁺			
Sr			

16. Fill-in the table below with proper quantum numbers to describe the last electron in each atom. Also, determine if the element is diamagnetic or paramagnetic. [3 points each]

Quantum #	In	Rh	Po	Mg
n				
l				
m_l				
m_s				

<i>dia- or para-</i>				
----------------------	--	--	--	--

17. Calculate the de Broglie wavelength in meters of a 45.9 g golf ball with a velocity of 52 m/s.
[3 points]

18. Explain the trends in atomic radius across a period (from left to right) and down a group (from top to bottom). [4 points]

19. Consider the electronic transition within a hydrogen atom from $n = 5$ to $n = 2$.
- Calculate the energy of the photon that is emitted in joules. [2 points]
 - Calculate the frequency in Hertz. [2 points]
 - Calculate the wavelength in meters. [2 points]
 - Does this emission occur within the visible region of the electromagnetic spectrum? [2 points]

20. Einstein won the Nobel Prize in 1922 for his explanation of the photoelectric effect where it was observed that many metals emit electrons when a light shines on their surface. (This is the basic idea behind solar panels.) It was expected that brighter lights would cause electron emission, while dim lights would not. However, experiments demonstrated that a high-frequency light from a dim source caused electron emission, while a bright low-frequency light did not. How did Einstein explain this unexpected result? (Feel free to include diagrams and equations to support your explanation.) [4 points]

Extra Credit: Of the 29 scientists who attended the 1927 Solvay Conference, 17 won the Nobel Prize. But only one, the lone woman to attend, won the Nobel Prize in two different fields. What was her name? [2 points]

**Formulas & Constants
(you may or may not need)**

1 inch = 2.54 cm (exact)

1 mile = 5280 ft \approx 1.609 km

1 kg \approx 2.205 lb

1 lb = 16 oz \approx 453.6 g

1 gal = 4 qt = 8 pt \approx 3.785 L

1 L = 1000 cm³

K = °C + 273.15

°F = 1.8 x °C + 32

°C = (°F – 32)/1.8

1 cal = 4.184 J

1 Cal = 1000 cal

q = m x C x ΔT

Avogadro's # = 6.022x10²³

Molar volume = 22.4 L

R = 0.08206 $\frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}}$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$u_{\text{rms}} = \sqrt{\frac{3RT}{M}}$$

$$KE = \frac{1}{2}mv^2 = \frac{3}{2}RT$$

1 atm = 760 mmHg

1 mmHg = 1 torr

P_{Total} = P₁ + P₂ + ...

P_A = X_A·P_{Total}

PV = nRT

ΔE = q + w

w = -PΔV

q = C x ΔT

q = m x s x ΔT

$$\Delta H^\circ_{\text{rxn}} = \Sigma[n \Delta H_f^\circ(\text{products})] - \Sigma[n \Delta H_f^\circ(\text{reactants})]$$

R = 8.314 J/mol·K

h = 6.626x10⁻³⁴ J·s

c = 2.9979x10⁸ m/s

R_H = 2.18x10⁻¹⁸ J

1 Hz = s⁻¹


$$\lambda = \frac{h}{mv}$$

$$\Delta x \times m\Delta v \geq \frac{h}{4\pi}$$

$$E_{\text{photon}} = h\nu = \frac{hc}{\lambda}$$

$$E_{\text{photon}} = R_H \left[\left(\frac{1}{n_{\text{final}}^2} \right) - \left(\frac{1}{n_{\text{initial}}^2} \right) \right]$$

The Electromagnetic Spectrum



Scratch Page
(to be handed in)